


Solar and Stellar Winds

The Solar Wind

Early evidence that the sun might be continuously expelling plasma at a high speed came from observations of the **dual tails** of comets.

One tail, made of **dust** slowly driven away from the comet by solar radiation, has an orientation that is tilted to the anti-sun (radial) direction by the comet's own orbital motion.

A second tail comes from cometary **ions** picked by the solar wind. It's more radial orientation implies that the radial outflow of the sun is much faster than the orbital speed of the comet.



The cause of the solar wind is the pressure expansion of the very hot (million degrees Kelvin) solar corona.


The high temperature causes the corona to emit X-rays.

Images made by orbiting X-ray telescopes show the solar corona has a high degree of spatial structure, organized by magnetic fields. Within **closed** field coronal **loops**, these effectively hold back the coronal expansion. But along radially oriented, open-field regions the wind flows rapidly outward, leading to a relative reduction of the plasma density that appears as a relatively dark "**coronal hole**".

Stan Owocki, Bartol Research Institute, University of Delaware

The Sun and other stars are commonly characterized by the **radiation** they emit.

But the past half-century has seen the discovery that the sun, and probably all stars, also **lose mass** through an essentially continuous, high-speed outflow or "**wind**".




The corona can also be observed in white light from the ground during a **solar eclipse**, or using "coronagraphs" with occulting disks that artificially eclipse the bright solar disk.

Such images show the closed loops are extended outward into radial **coronal streamers** by the wind outflow.


Both X-ray and white-light observations show that closed-field **loops** tend to occur near the **equator**, while open-field coronal **holes** are usually near the solar **poles**.

But the solar wind is most directly observed *in situ* by an interplanetary spacecraft with plasma instruments to measure the wind's **speed**, elemental composition, ionization state, and the interplanetary magnetic field (**IMF**).



The generally lower-speed ecliptic-plane wind also shows abrupt switches to high-speed streams that originate from low-latitude coronal holes.

The rotation of the sun brings about a collision between these high- and low-speed streams along spiral **Co-rotating Interaction Regions**, forming abrupt shock discontinuities in plasma conditions that are measured by spacecraft, often with a repetition close to the solar rotation period.



The solar wind interacts with the earth's **magnetosphere**, providing a key way that solar activity can induce geomagnetic activity, and perhaps even influence earth's climate and weather.


Finally, the solar wind blows out a "**heliospheric cavity**" in the local interstellar medium. The Voyager spacecraft may reach the "**bow shock**" of this cavity within the next couple decades.

Stellar Winds

Evidence of episodic stellar mass loss in the form of novae or supernovae has been known since antiquity. But the realization that star's could also have **continuous** wind dates from the 1960's, largely from analogy with the solar wind.

Low-density, **optically thin** coronal winds from solar-like, low mass, main-sequence stars can only be inferred indirectly, e.g. by X-ray observations suggesting stellar coronae.

But for some stars -- e.g. during the Red Giant phase of a solar-mass star, or from hot, luminous, high-mass stars -- the stellar winds are dense enough to be **optically thick** in spectral lines.




Lines formed by scattering of the stellar radiation with the expanding wind develop a characteristic shape -- a **P-Cygni** profile -- whose features provide a direct diagnostic of key wind parameters, like the wind **speed** and **mass loss rate**.

For **Red Giant** stars, such profiles suggest relatively **slow** speeds, 10-50 km/s, but with mass loss rates up to **million** times that of the solar wind, i.e., $\sim 10^{-8} M_{\odot}/\text{yr}$.

But massive stars show the strongest winds, with speeds sometimes exceeding 3000 km/s, and mass loss rates up to a **billion** times the solar wind, i.e. $\sim 10^{-5} M_{\odot}/\text{yr}$!


This is large enough that, during the course of their relatively brief ($\sim 10^7$ yr) evolutionary lifetime, such massive stars can be stripped of their entire hydrogen envelope, exposing a "**Wolf-Rayet**" star characterized by strong line emission from ions of nuclear processed elements like Carbon, Nitrogen, and Oxygen.

For **Red Giants** the wind **driving** mechanism is **not** well understood. but may involve a combination of **radiation pressure** on dust.



For hot, luminous stars the driving is generally thought to stem from **radiation pressure** acting through **line** scattering. The **Doppler shift** of the line-profile within the expanding wind effectively "sweeps out" the star's continuum momentum flux..




This makes the driving **force** a function of the wind velocity and **acceleration**, leading to strong **instabilities** that likely make such winds highly **turbulent**.



Monitoring campaigns of P-Cygni lines formed in hot-star winds also often show **modulation** at periods comparable to the stellar rotation period.

These may stem from large-scale surface structure that induces wind variation analogous to solar **Corotating Interaction Regions**.

The generally rapid rotation of hot stars can also lead to focusing of the outflow into an **an equatorial disk**.



The large mass loss of hot-stars also represents a substantial **source** of energy and **mass** into the **interstellar medium**.

Indeed, interstellar nebulae near young star clusters often show clear "**wind-blown bubbles**" from the many hot, massive stars.

In particularly dense clusters, these can even coalesce into large "**superbubbles**".



The compression around such wind bubbles may play a role in **triggering** further **star formation**. Some galaxies even appear to be undergoing "**starbursts**", with integrated spectra dominated by young, massive stars.

Radiative driving processes similar to those occurring in hot-star winds may even be key to understanding outflows from **Active Galactic Nuclei** and **quasars**.

